

Multimodal CEA-targeted image-guided colorectal cancer surgery using ¹¹¹In-labeled SGM-101

Jan Marie de Gooyer^{1,2}, Fortuné M. K. Elekonawo^{1,2}, Desirée L. Bos¹, Rachel S. van
 der Post³, André Pèlegri⁴, Bérénice Framery⁵, Françoise Cailler⁵, Alexander L.
 Vahrmeijer⁶, Johannes H. W. de Wilt², Mark Rijpkema¹

¹Radboud university medical center, department of Radiology, Nuclear Medicine & Anatomy, Nijmegen, the
 Netherlands. ²Radboud university medical center, department of Surgical oncology, Nijmegen, the Netherlands.
³Radboud university medical center, department of Pathology, Nijmegen, the Netherlands. ⁴IRCM, Institut de
 Recherche en Cancérologie de Montpellier, INSERM U1194, Université de Montpellier, Institut régional du
 Cancer de Montpellier, Montpellier, F-34298, France. ⁵SurgiMab, Montpellier, France. ⁶Leiden university medical
 center, department of Surgery, Leiden, the Netherlands.

Running title: Multimodal image-guided colorectal cancer surgery

Key words: Colorectal cancer, Image guided surgery, fluorescence imaging, preoperative imaging,
 multi-modal imaging

Corresponding author

Jan-Marie de Gooyer
 Radboudumc, Geert Grooteplein Zuid 10
 6525 GA Nijmegen
 +31 (0)24 366 72 44
 jan-marie.gooyer@radboudumc.nl

Disclosure statement

FC and BF are employed by Surgimab, and are inventors on the patent that protects SGM-101. FC, BF
 and AP are stockholders of Surgimab. All other authors declare no competing interests.

Translational relevance statement

With surgical procedures for colorectal cancer becoming more complicated and extensive, adequate localization and delineation of tumors can be arduous for surgeons. Intraoperative dual-modality image guided surgery with multimodal tracers is a powerful technique that may provide essential information about tumor margins and the location of metastatic lesions. In the current study, we report on the synthesis, *in vitro* characterization, *in vivo* testing and translational evaluation of a novel multimodal CEA-targeting tracer for image guided colorectal cancer surgery. SPECT/CT and fluorescence imaging clearly visualized specific tracer uptake in a colorectal peritoneal carcinomatosis model, which was confirmed by immunohistochemical analysis and gamma counting. The translational *ex vivo* incubation study with fresh surgically derived human peritoneal colorectal cancer deposits showed high tumor to background ratios of the fluorescence and radiosignal. These results demonstrate the clinical potential our multimodal CEA-targeting tracer for image guided surgery and facilitate rapid clinical implementation, which is currently in progress.

Abstract

Purpose: Intraoperative image guidance may aid in clinical decision making during surgical treatment of colorectal cancer. We developed the dual-labeled CEA-targeting tracer [^{111}In]In-DTPA-SGM-101 for pre- and intraoperative imaging of colorectal cancer. Subsequently we investigated the tracer in preclinical biodistribution and multimodal image-guided surgery studies, and assessed the clinical feasibility on patient derived colorectal cancer samples, paving the way for rapid clinical translation.

Experimental design: SGM-101 was conjugated with p-isothiocyanatobenzyl (ITC)-diethylenetriaminepentaacetic acid (DTPA) and labeled with ^{111}In . The biodistribution of 3, 10, 30, and 100 μg [^{111}In]In-DTPA-SGM-101 was assessed in a dose escalation study in BALB/c nude mice with subcutaneous LS174T human colonic tumors followed by a study to determine the optimal time point for imaging. Mice with intraperitoneal LS174T tumors underwent microSPECT/CT imaging and fluorescence image-guided resection. In a final translational experiment, we incubated freshly resected human tumor specimens with the tracer and assessed the tumor-to-adjacent tissue ratio of both signals.

Results: The optimal protein dose of [^{111}In]In-DTPA-SGM-101 was 30 μg (tumor-to-blood ratio 5.8 ± 1.1) and the optimal timepoint for imaging was 72 hours after injection (tumor-to-blood ratio 5.1 ± 1.0). In mice with intraperitoneal tumors, [^{111}In]In-DTPA-SGM-101 enabled preoperative SPECT/CT imaging and fluorescence image-guided resection. After incubation of human tumor samples overall fluorescence and radiosignal intensities were higher in tumor areas compared to adjacent non-tumor tissue ($p < 0.001$).

Conclusions: [^{111}In]In-DTPA-SGM-101 showed specific accumulation in colorectal tumors, and enabled micro-SPECT/CT imaging and fluorescence-guided tumor resection. Thus, [^{111}In]In-DTPA-SGM-101 could be a valuable tool for preoperative SPECT/CT imaging and intraoperative radioguided localization and fluorescence-guided resection of colorectal cancer.

Introduction

Colorectal cancer is currently the second leading cause of cancer death in women, and the third most common cause in men(1). While therapeutic strategies have evolved throughout the last decades, radical surgical resection of all tumor tissue with clear margins (R0) remains the mainstay of treatment. Surgical treatment has become increasingly versatile with procedures such as cytoreductive surgery for colorectal peritoneal metastasis, or extensive exenterative surgery for locally advanced or recurrent rectal cancer(2, 3). Complete tumor resection is pivotal for successful treatment outcome, but remains challenging in many patients. In rectal cancer, involvement of the circumferential resection margin (CRM) still occurs in 1 to 28% of all surgically treated cases(4). There are several factors that can complicate surgical treatment, such as distorted pelvic anatomy after previous resections, multifocal tumor deposits, the presence of multiple small tumor nodules in peritoneally metastasized cancers, and difficulty to distinguish benign from malignant tissue, for example after neo-adjuvant therapy(5). Real-time intraoperative margin assessment and tumor visualization techniques guiding the surgeon may be advantageous to overcome these obstacles.

Carcinoembryonic antigen(CEA)-targeted fluorescence imaging is a technique that can be of added value in surgical treatment of both locoregional and metastasized colorectal cancer. CEA is a widely known glycoprotein and tumor marker that is highly expressed on almost all (more than 95%) colorectal cancers, but also on a variety of other malignancies such as gastric, pancreatic, lung, breast and esophageal cancer(6). SGM-101 is a CEA-specific chimeric antibody conjugated to a near-infrared fluorophore that has been developed to facilitate CEA-targeted fluorescence image-guided surgery of colorectal cancer(7).

Preclinical toxicology, pharmacology and pharmacokinetic results have demonstrated the absence of significant adverse effects of SGM-101 at doses well above the anticipated maximal human exposure (8). A previous phase I/II clinical trial has also already shown the potential of SGM-101 for intraoperative detection of colorectal cancer tissue. In this trial, CEA-targeted fluorescence-guided

surgery resulted in the detection of additional malignant lesions that were not clinically suspect, emphasizing the potential of SGM-101 for intraoperative tumor detection of colorectal cancer(9).

While the use of targeted near-infrared fluorescence-guided surgery is of great value to enhance oncological procedures, the limited tissue penetration depth of NIR-fluorescent light confines application to superficial tumors. Accurate visualization of deeper located tumors is difficult and only feasible after proper surgical exposure(9). Adding a radiolabel to a fluorescent tumor-targeting tracer may overcome this limitation by facilitating intraoperative radioguidance(10). Such a dual-labeled tracer also enables preoperative imaging with SPECT/CT and straightforward quantification in (pre)clinical research by gamma counting(11). Quantitative assessment of tracer accumulation and subsequent determination of the tumor to background ratio is a key determining factor of tumor detectability and sensitivity of the imaging technique, but accurate quantification of NIR-fluorescence is difficult and reported values differ widely(12). Thus, a tracer containing both a fluorophore and a gamma radiation emitting radionuclide may be a powerful approach(13).

In this study, SGM-101 was conjugated to a radiolabel (^{111}In) to create the dual-labeled tracer [^{111}In]In-DTPA-SGM-101. We evaluated the characteristics, immunoreactivity and stability of this new tracer in vitro. Subsequently we determined the in vivo biodistribution, and performed CEA-targeted micro-SPECT and fluorescence image-guided surgery in a CEA-expressing colorectal cancer xenograft model. In a final translational experiment, we investigated this new tracer in an ex vivo incubation experiment with patient derived surgically resected peritoneal colorectal cancer deposits.

Materials & Methods

Synthesis of DTPA-SGM-101

SGM-101 is a highly specific anti-CEA chimeric monoclonal antibody(ch511) conjugated to a fluorophore in the 700 nm range (BM104). Clinical grade SGM-101 was obtained from Surgimab (Montpellier, France)(7). First, SGM-101 was dialyzed for 5 days in a Slide-A-Lyzer (10-kDa cutoff; Pierce Biotechnology Inc, Waltham, MA, USA.) against a 0.5M phosphate-buffered saline (PBS) solution. Subsequently, SCN-Bz-DTPA (Macrocyclics, Plano, TX, USA) was added in a 3-fold molar excess in Dimethyl sulfoxide (DMSO). After 1 hour of incubation at room temperature on an orbital shaker in the dark, the reaction mixture was dialyzed for 5 days in a Slide-A-Lyzer (10-kDa cutoff; Pierce Biotechnology Inc, Waltham, MA, USA.). against PBS. The final concentration of the new conjugate was determined spectrophotometrically at 280 nm using an Infinite M200 Pro multimode reader (Tecan group Ltd, Männedorf, Switzerland). The optical properties of both the SGM-101 starting material and the new conjugate were investigated. Absorbance and fluorescence spectra of both solutions were determined using the Infinite M200 Pro multimode reader. A radiolabeling of the new DTPA-SGM-101 with Indium-111 was done (see below) before and after dialysis to calculate the conjugation ratio of the chelator DTPA to SGM-101. HPLC analysis of the radiolabeled conjugate was performed to determine the radiochemical purity of the new compound.

Radiolabeling and Quality Control of DTPA-SGM-101

For the first 2 biodistribution experiments on mice with subcutaneous tumors, DTPA-SGM-101 was labelled with 0.5 MBq of ¹¹¹In (Curium, Petten, the Netherlands) in 0.5M 2-(N-morpholino) ethanesulfonic acid (MES) buffer, pH 5.4 (2 times the volume of the [¹¹¹In]InCl₃ solution), and incubated for 20 min at room temperature under metal-free conditions. For studies where SPECT/CT was performed, DTPA-SGM-101 was radiolabeled with 10 MBq of ¹¹¹In per mice under similar conditions. Subsequently, 50 ethylenediamine tetraacetic acid (EDTA) solution was added (final concentration, 5 mM) to chelate unincorporated ¹¹¹In. Final labelling efficiency of the compound was

determined by instant thin-layer chromatography on silica gel strips (Agilent Technologies, Amstelveen, The Netherlands), where a 0.15 M citrate buffer with pH 6.0 was used as mobile phase. The labelling efficiency exceeded 98% for all experiments. The immunoreactivity of [¹¹¹In]In-DTPA-SGM-101 was determined with LS174T cells, as described by Lindmo et al(14). In short: A serial dilution of LS174T cells was incubated with [¹¹¹In]In-DTPA-SGM-101 for 30 min at 37°C. Subsequently, cells were centrifuged and analyzed in a shielded 3"-well-type γ-counter (Perkin-Elmer, Boston, MA, USA). The LS174T colorectal cancer cells used in all experiments were originally acquired from the American Type Culture Collection (ATCC, Manassas, VA, USA, Batch F-9126). Cells were last tested for *Myxoplasma* in January 2018. All experiments were conducted within 1 to 2 months after thawing.

Tumor models

All animal experiments were conducted with 6- to 8-week-old female athymic BALB/c nu/nu mice (Janvier, Le Genest-Saint-Isle, France) weighing 20–25 g. Mice were given at least 1 week to become accustomed to laboratory conditions before starting with the experiments. Mice were housed in individually ventilated filter topped cages under nonsterile standard conditions with free access to chlorophyll free chow and water in accordance with institutional guidelines. The subcutaneous tumors were induced by subcutaneous injection of 3×10^5 LS174T cells in 200 μL of RPMI 1640 medium (Gibco, Thermo Fischer Scientific, Waltham, MA, USA) in the right flank of the mice. Intraperitoneal tumors were induced as described previously by Koppe et al(15). In summary, mice were injected intraperitoneally with 10^6 LS174T cells suspended in 300 μL of RPMI 1640 medium. Experiments were continued 1-2 weeks after tumor cell injection.

SPECT/CT and Fluorescence imaging

SPECT/CT and fluorescence imaging was performed as described previously by Rijpkema et al(16). In summary, SPECT/CT images were acquired with the U-SPECT-IICT system (MILabs, Utrecht, the

Netherlands). Mice were imaged for 2 frames of 25 minutes using a 1.0-mm-diameter pinhole collimator, followed by a CT scan for anatomic referencing (spatial resolution 160 μ m; 65 kV; 612 μ A). All scans were reconstructed with software supplied by MILabs, using an ordered subset expectation maximization algorithm with a voxel size of 0.375 mm. SPECT/CT images were subsequently analyzed and created using the Inveon Research Workplace Software (IRW, version 4.1). Fluorescence images were acquired with an in vivo imaging system (Xenogen VivoVision IVIS Lumina II; Caliper Life Sciences, Hopkinton, MA, US). The acquisition time was 1–2 min with the following parameters: F/stop 2; binning medium; field of view C; excitation filter 605 nm; emission filter 695–770 nm. Correction for autofluorescence was performed using a second acquisition of 10 s with an excitation filter of 605 nm.

Immunohistochemistry

Immunohistochemistry was performed as described by Elekonawo et al (17). Slides were deparaffinized with xylene, rehydrated in ethanol and rinsed in distilled water according to standard local protocol. Heat-induced antigen retrieval was performed in EDTA solution (pH 9.0). Endogenous peroxidase activity was blocked with 3% H₂O₂ in 10 mM PBS for 10 minutes at room temperature. Subsequently, tissue sections were washed with 10 mM PBS and stained with primary antibodies (CEA: Mouse monoclonal; Clone COL-1; MS-613-P; Neomarkers Fremont, CA, USA). Next, sections were incubated with biotin free Poly-HRP-anti mouse/rabbit IgG (ImmunoLogic/VWR International B.V., Amsterdam, The Netherlands) in EnVision™ FLEX Wash Buffer (Dako Denmark A/S, Glostrup, Denmark) (1:1) at room temperature for 30 minutes. Antibody binding was visualized using the EnVision™ FLEX Working Solution (Dako Denmark A/S, Glostrup, Denmark) at room temperature for 10 minutes. All sections were counterstained with hematoxylin for 5 seconds and subsequently dehydrated in ethanol and coverslipped.

Ethical Approval

All animal experiments were approved by the Animal Welfare Committee of the Radboud University Medical Center (Local code: 2015-0071-147) and were conducted in accordance with the Revised Dutch Act on Animal Experimentation (Wet op de Dierproeven). The experiments involving the use of surgically derived human tissue samples have been carried out according to the principles of the Declaration of Helsinki. The study protocol and collection of tissue were approved by the research ethics committee of the region Arnhem-Nijmegen (CMO regio Arnhem-Nijmegen, file code 2019-5437).

In Vivo Experiments

Dose escalation study

Mice with subcutaneous LS174T tumors on their right flank received an intravenous injection of 3, 10, 30, or 100 µg of [¹¹¹In]In-DTPA-SGM-101 per mouse (5 mice per group). A 5th group received 30 µg of radiolabeled [¹¹¹In]In-DTPA-SGM-101 complemented with 1 mg of unlabeled DTPA-SGM-101 as control group. Three days after injection of [¹¹¹In]In-DTPA-SGM-101, mice were euthanized using O₂/CO₂ asphyxiation and blood, lung, heart, spleen, pancreas, stomach, small intestine, kidney, liver, muscle and tumor were dissected and weighed. Activity in all resected specimens was measured in a gamma-counter (2480 WIZARD2; PerkinElmer). To calculate the uptake of radiolabeled antibodies in each sample as a fraction of the injected activity, aliquots of the injected dose were counted simultaneously. The results were expressed as percentage injected activity per gram (%IA/g).

Optimal time point determination

Mice with subcutaneous LS174T tumors received the optimal protein dose of 30 µg (as determined in the previous dose escalation study) of [¹¹¹In]In-DTPA-SGM-101 (10 MBq per mouse). After 24, 48, or 72 hours (5 mice at each time point), SPECT/CT and fluorescence imaging was performed as described above. Subsequently, mice were euthanized and the biodistribution of the radiolabel was

determined as described in the previous experiment to assess the optimal timepoint for imaging after administration of the dual-labeled tracer.

Preoperative SPECT and fluorescence image-guided surgery study

Using the optimal conditions as determined in the first two in vivo studies, multimodal image-guided surgery was performed in mice with intraperitoneally growing LS174T tumors as previously described by Rijpkema et al (16). Mice received 30 µg of [¹¹¹In]In-DTPA-SGM-101 (10 MBq per mouse; 4 mice) or 30 µg of [¹¹¹In]In-DTPA-SGM-101 (10MBq per mouse, 2 mice) with 1 mg of unlabeled DTPA-SGM-101 that served as a negative control. Three days after administration of the antibody conjugate, preoperative SPECT/CT images were acquired, followed by fluorescence imaging (as described above) of the mice in supine position after surgical relocation of skin, abdominal muscle layers, and peritoneum. Subsequently, the visualized tumor nodules were resected and fluorescence imaging was repeated to check for residual tumor tissue. This procedure was repeated until all tumor nodules were removed.

Ex vivo incubation of fresh patient-derived peritoneal tumor samples

Finally, a translational experiment was conducted to assess tumor to surrounding (normal) tissue contrast in human tissue samples using fluorescence imaging and autoradiography as previously described by Elekonawo et al(17). Fresh surgically obtained peritoneal colorectal cancer deposits were incubated in tissue culture medium containing the tracer [¹¹¹In]In-DTPA-SGM-101. Immediately after surgical resection, colorectal peritoneal tumor deposits were acquired and subsequently incubated overnight at 37°C in DMEM supplemented with 1% penicillin/streptomycin and 0.1% bovine serum albumin (BSA)(Sigma-Aldrich Chemie N.V., Zwijndrecht, Netherlands) and 4 µg·ml⁻¹, [¹¹¹In]In-DTPA-SGM-101 (0.05 MBq/ug). After incubation overnight, tumor deposits were washed in at least 3L of continuously moving serially refreshed phosphate-buffered saline supplemented with 0.1% BSA at 4-8°C for a minimum of 6 hours. Specimens were subsequently fixed in formalin and

embedded in paraffin. Four μm tissue sections were cut and placed on glass slides. Fluorescence images were acquired on a closed field fluorescence imaging system (Odyssey CLx; LI-COR Biosciences, Lincoln, NE, USA). Slides were then placed on a photostimulable phosphor plate. After 7 days, we imaged the plate on a phosphor imager (Typhoon FLA 7000 phosphor imager, GE Healthcare, Hoevelaken, The Netherlands) Subsequently slides were stained with eosin & haematoxylin (H&E) and immunohistochemistry (IHC) of carcinoembryonic antigen (CEA)(mouse monoclonal; Clone COL-1; MS-613-P; Neomarkers Fremont, CA, USA) was performed. Surgical specimens from two patients received a non-specific hIgG antibody-conjugate (SouthernBiotech, Birmingham, AL, USA) as negative controls. An independent pathologist reviewed the H&E and CEA slides to verify the presence and location of tumor cells. Regions of interest (ROI) (tumor and adjacent normal tissue) were drawn in the corresponding fluorescence and autoradiography images to determine tumor to background ratios of both fluorescence and radiosignal.

Statistical analysis

Statistical analyses were performed using SPSS Statistics 25.0 (IBM) and GraphPad Prism software (version 5.03; GraphPad Software). Tissue uptake at the different protein dose levels and time points was tested for significance using a 1-way ANOVA test with post hoc Bonferroni correction. To test for differences in tumor and normal fluorescence and autoradiography intensities, a Mann-Whitney U test was used. Mean tumor to background ratios of the experimental group were compared to mean tumor to background ratios in the control groups with an independent samples t-test. An α -value < 0.05 was considered to be statistically significant in all analyses.

Results

Comparison of SGM-101 and DTPA-SGM-101 showed similar patterns of absorbance and fluorescence emission before and after conjugation to the chelator (supplemental figure 1). The substitution ratio of SGM-101 and DTPA was 1. The immunoreactive fraction of all conjugates

exceeded 80% and radiochemical purity of [^{111}In]In-DTPA-SGM-101 used in all studies was >99%.

In the dose escalation study, [^{111}In]In-DTPA-SGM-101 specifically accumulated in the subcutaneous CEA-positive LS174T tumors, with mean tumor uptake of 56.3 ± 14.2 , 42.5 ± 15.1 , 46.2 ± 6.7 and 21 ± 1 %IA/g after injection of 3, 10, 30 and 100 μg , respectively. Tumor to blood ratios of mice that were injected with 3, 10, 30 or 100 μg were 5.6 ± 1.3 , 5.0 ± 0.9 , 5.8 ± 1.1 and 2.3 ± 0.4 , respectively. Adding an excess of 1 mg unlabeled DTPA-SGM-101 decreased tumor uptake to 5.3 ± 0.4 %IA/g and the tumor-to-blood ratio to 0.53 ± 0.2 %IA/g, illustrating that tumor uptake of [^{111}In]In-DTPA-SGM-101 was specific. Based on these results, 30 μg was chosen as the optimal dose for subsequent experiments. Results of the biodistribution are summarized in figure 1.

In the second study, SPECT/CT imaging, NIR-fluorescence imaging and biodistribution studies were performed 24, 48 and 72 hours after administration. Tumor uptake was 34.5 ± 5.4 , 42.0 ± 16.1 and 56.4 ± 13.9 %IA/g after 24, 48 and 72 hours respectively. Tumor to blood ratios were 2.5 ± 0.5 after 24 hours, 3.7 ± 0.8 after 48 hours and 5.1 ± 1.0 after 72 hours. Tumors could be visualized with both SPECT and fluorescence imaging at each time point. Seventy-two hours after administration of [^{111}In]In-DTPA-SGM-101 was chosen as the optimal time point for the imaging of peritoneal tumors in the next experiment. The results of this study determining the optimal timepoint are summarized in supplemental figure 2.

In the third in vivo experiment, we performed preoperative SPECT/CT imaging and fluorescence-guided resection of tumor deposits in mice with intraperitoneally growing tumors. First, SPECT/CT images were acquired to visualize the extent of peritoneal disease dissemination. Subsequently mice were euthanized, and laparotomy was performed to search for tumor deposits. After suspected tumors were found by combining visual inspection and SPECT/CT imaging, fluorescence images were acquired and tumors were resected. Figure 2 shows that after resection of all macroscopically visible

tumors, fluorescence imaging revealed submillimeter tumor deposits that were initially missed by visual inspection. Mice that received the optimal dose with the addition of 1 mg unlabeled DTPA-SGM-101 did not show any specific accumulation of the tracer in the tumor on both imaging modalities (fluorescence imaging and SPECT). Immunohistochemical analysis showed clear concordance between CEA-expressing tumor cells, fluorescence and radiosignal (Supplemental figure 3).

In a final translational experiment, we collected 43 peritoneal deposits from 11 patients (3 male, 8 female). Median age was 76 (65-78) years. Seven patients were diagnosed with a peritoneally metastasized adenocarcinoma. In one patient the peritoneal metastases originated from a mucinous adenocarcinoma and in another patient from a signet-ring cell carcinoma. In one case the peritoneal deposits contained a low grade appendiceal mucinous neoplasm (LAMN), and one other case was an epithelial ovarian cancer. These last two cases did not express CEA and were used as negative controls.

After the incubation experiment, subsequent series of slides were produced from the surgically resected tumor deposits. This resulted in 43 series of slides that were analyzed. Thirty-three of the resected peritoneal deposits contained malignant cells. The other 10 lesions did not contain vital malignant tumor cells and consisted of fibrosis, necrosis, mucin, stroma or fat. Median fluorescence intensity was 1371 (IQR: 1020-1834) in tumors. This was significantly higher compared to adjacent normal tissue (fat, peritoneum or stroma) where the median fluorescence intensity was 293 (IQR: 168 - 317) ($p < 0.001$). Median intensity of the autoradiography for tumor tissue was $10.46 \cdot 10^6$ (IQR: $5.192 \cdot 10^6 - 14.66 \cdot 10^6$). The median autoradiography intensity in adjacent non-tumor tissue was $1.913 \cdot 10^6$ (IQR: $1.142 \cdot 10^6 - 2.724 \cdot 10^6$) ($p < 0.001$). Tumor to background ratios for the fluorescence and radiosignal of all cases are shown in figure 3. Fluorescent and radiosignal-based tumor to background ratios are significantly lower in the CEA-negative cancers (no radiography data available for the LAMN due to a technical failure) and in the control cases incubated with the aspecific hIgG (p

< 0.001 for both signals). Three examples of slides (H&E, CEA, fluorescence and autoradiography) are provided in figure 4, illustrating the concordance between CEA-positive tumor cells and fluorescence and radiosignal. Microscopic images at a higher magnification confirming colocalization of CEA-expressing tumor cells and the fluorescence signal are shown in supplemental figure 4.

Discussion

This study shows the in vitro and in vivo characterization of [^{111}In]In-DTPA-SGM-101 and the feasibility of multimodal image-guided surgery using this tracer. Accumulation of [^{111}In]In-DTPA-SGM-101 was high and specific in subcutaneous and peritoneal CEA-expressing LS174T xenografts. The biodistribution of this ^{111}In -labeled fluorescent tracer showed high tumor to blood ratios and low uptake in other organs. Translational ex vivo incubation studies with fresh surgically derived human peritoneal colorectal cancer deposits showed high tumor to background ratios and overall correspondence of both the fluorescence and radiosignal, suggesting the feasibility of this tracer for multimodal image-guided surgery of CEA-expressing colorectal cancer, warranting further evaluation in clinical studies.

The biodistribution of SGM-101 has been extensively investigated by Gutowski et al. in a similar colorectal cancer model(7). The reported results show a highly similar in vivo distribution of the tracer, with high uptake in the tumor compared to other tissues. As expected, conjugation of DTPA to SGM-101 and subsequent radiolabeling with ^{111}In did not affect the in vivo behavior and distribution of SGM-101, as in accordance with the literature(18-20). The substitution ratio of DTPA to SGM-101 in our study was 1, which proved sufficient for radiolabeling in all experiments. Most chelating molecules can be conjugated to full IgG antibodies in ratios of up to 5 without significantly influencing or altering the immunoreactivity and in vivo behavior of the antibody(18-20). Since the in vivo behavior of iodinated SGM-101 and [^{111}In]In-DTPA-SGM-101 is comparable, [^{111}In]In-DTPA-SGM-101 could be used for quantification of tracer in tissue and blood, facilitating pharmacokinetic analysis of SGM-101 in a clinical setting.

In addition to accurate tracer quantification and pharmacokinetic analysis, intravenous injection of this dual-labeled tracer could facilitate clinical decision making by improving preoperative tumor imaging. Whole body SPECT/CT imaging might be used as an additional tool for non-invasive preoperative staging of the primary tumor, distant metastases and lymph node metastases in colorectal cancer patients, potentially adding to improved selection of patients with peritoneal metastases who are eligible for cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. Conventional imaging modalities tend to underestimate the degree of intraperitoneal tumor extent in these patients(21). As a result of this underestimation, surgical procedures are performed on patients that are diagnosed with irresectable disease after surgical exploration of the abdomen, resulting in futile laparotomies or incomplete cytoreduction(21-24). SGM-101 fluorescence-guided cytoreductive surgery for colorectal peritoneal carcinomatosis has already been tested in a multicenter pilot study and in one third of all cases, detection of additional malignant lesions by fluorescence imaging led to an adjustment of the peritoneal carcinomatosis index (PCI) (25). Preoperative SPECT/CT imaging might also be used as a diagnostic aid to determine whether patients have extra-abdominal disease. Current criteria for determining if lymph nodes or lung nodules are malignant are mainly based on size and morphological features. A tumor-targeted molecular imaging approach might be valuable in determining whether lymph nodes and lung nodules that are considered borderline on conventional CT are malignant or benign. Combining these benefits of fluorescence-guided surgery with the additional value of a radiolabel could potentially improve diagnosis and treatment of these patients even further and will be evaluated in a clinical trial.

Another potential application of preoperative SPECT/CT imaging is the evaluation of metastatic lymph node status in patients with rectal cancer after preoperative radiation based treatment. The current imaging modality of choice for determining the lymph node status is Magnetic Resonance Imaging (MRI), but reported sensitivity is low(26). CEA targeted preoperative SPECT/CT could potentially be used as a restaging modality after neoadjuvant (chemo)radiotherapy since CEA-

expression is not influenced by neoadjuvant treatment(27). Subsequently, if patients exhibit tracer uptake in lymph nodes outside the standard total mesorectal resection specimen, such as lateral pelvic lymph nodes, radioguidance with a gamma probe could perhaps aid in localizing these lymph nodes. Subsequent intraoperative real time fluorescence imaging could also help the surgeon in performing a complete resection of all metastatic lymph nodes and the primary tumor.

Besides colorectal cancer, combined CEA-targeted radioguidance and fluorescence imaging can potentially be translated to other CEA-expressing malignancies as well. De Gouw et al. have shown that 68% of lymph node metastases of esophageal cancer express CEA(28). Approximately one third of all surgically treated patients with esophageal cancer have lymph node metastases after neoadjuvant treatment, but lymphadenectomy is always performed because adequate treatment response assessment of lymph node metastases is not possible with current imaging modalities(29). CEA-targeted multimodal imaging might aid in minimizing or possibly omitting this procedure and reduce the associated morbidity in selected cases. The same hypothesis is worth investigating in gastric cancer, where similar dilemmas exist(30). CEA-targeted multimodal imaging could be investigated as a potential aid in identifying additional metastatic lymph nodes outside of the standard resection plane. It might also guide the surgeon in deciding to perform a less invasive D1 lymph node dissection instead of the more extensive D2 dissection which is associated with increased morbidity(31, 32). In pancreatic cancer, fluorescence-guided surgery with SGM-101 was successful despite suboptimal intrinsic characteristics such as poor vascularization, abundant presence of desmoplastic stroma and localization of solitary ducts of tumor cells within benign pancreatic tissue(33). However, tumor to background ratios were lower as compared to colorectal cancer(9, 33). The addition of a radiolabel could be of added value when performing CEA-targeted image-guided surgery in these patients, similar to Hekman et al, who successfully translated this approach of multimodal tumor-targeted image-guided surgery into the clinic(10, 34). In a first clinical study they demonstrated the feasibility of combining preoperative SPECT/CT imaging, intraoperative

gamma probe detection and NIR-fluorescence-guided surgery in patients with clear cell renal cell carcinoma(10).

There are some limitations to this study that have to be addressed. First, a mouse model with human colorectal cancer xenografts was used and one should consider that CEA-expression may be more heterogeneous in a clinical setting. Since we did not use transgenic human CEA-expressing mouse models, tumor to background ratios will not be directly translatable to the clinical setting. However, in the preclinical toxicology study, the biodistribution of iodinated SGM-101 did not differ significantly between transgenic CEA-expressing mice and wild type animals because physiological CEA-expression is exclusively located on the apical side of healthy epithelial cells, which is also the case with intestinal CEA-expression in humans (8, 35). Therefore it is not accessible to circulating CEA targeting molecules, as opposed to malignant colorectal tumors, where CEA is overexpressed on the entire cell surface at a 60-fold higher density (35, 36). Second, the translational experiment that was performed with surgically derived peritoneal tumor deposits relies on passive diffusion of the tracer into the tissue to establish adequate uptake and tumor to background ratios. Since passive diffusion of full IgG antibodies will only facilitate penetration of approximately 100 μm within 12 to 18 hours of incubation, the center of the tumor deposit may not be reached by the tracer (17). However, although tracer distribution after intravenous injection still has to be shown in the clinic, the edges and invasive margins of the tumor are the main regions of interest in surgical oncology and the current results show that sufficient tumor-to-background ratios were reached for both the radio- and the fluorescence signal. These results are comparable to the tumor to background ratios found in the clinical studies conducted with SGM-101 (9, 25, 33).

In conclusion, this study shows the feasibility of CEA-targeted multimodal SPECT/CT and fluorescence imaging using [^{111}In]In-DTPA-SGM-101 in colorectal cancer models. [^{111}In]In-DTPA-SGM-101 enables specific and sensitive detection of colorectal cancer lesions in vivo and specifically targets tumor lesions in fresh surgical samples of colorectal cancer. The addition of a gamma-emitting radiolabel to

SGM-101 enables straightforward quantification in vivo and may facilitate preoperative detection of tumors by SPECT/CT imaging. Subsequent radioguided intraoperative localization of deeper-seated lesions combined with CEA-targeted fluorescence guided surgery might further benefit the surgical treatment of colorectal cancer and other CEA-expressing malignancies. These promising preclinical results and our experience with the fluorescent tracer in clinical trials will pave the way for rapid clinical translation. Based on our results, this new multimodal tracer will likely improve clinical decision making by personalized precision surgery.

Acknowledgements

The authors would like to thank Bianca Lemmers van de Weem and Karin de Haas-Cremers for the assistance with the animal experiments.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*. 2018;68(6):394-424.
- Verwaal VJ, van Ruth S, de Bree E, van Sloothen GW, van Tinteren H, Boot H, et al. Randomized trial of cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy and palliative surgery in patients with peritoneal carcinomatosis of colorectal cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2003;21(20):3737-43.
- Brown KGM, Solomon MJ, Koh CE. Pelvic Exenteration Surgery: The Evolution of Radical Surgical Techniques for Advanced and Recurrent Pelvic Malignancy. *Dis Colon Rectum*. 2017;60(7):745-54.
- Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2008;26(2):303-12.
- Dresen RC, Kusters M, Daniels-Gooszen AW, Cappendijk VC, Nieuwenhuijzen GA, Kessels AG, et al. Absence of tumor invasion into pelvic structures in locally recurrent rectal cancer: prediction with preoperative MR imaging. *Radiology*. 2010;256(1):143-50.
- Hammarström S. The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues. *Semin Cancer Biol*. 1999;9(2):67-81.
- Gutowski M, Framery B, Boonstra MC, Garambois V, Quenet F, Dumas K, et al. SGM-101: An innovative near-infrared dye-antibody conjugate that targets CEA for fluorescence-guided surgery. *Surg Oncol*. 2017;26(2):153-62.

8. Framery B, Gutowski M, Dumas K, Evrard A, Muller N, Dubois V, et al. Toxicity and pharmacokinetic profile of SGM-101, a fluorescent anti-CEA chimeric antibody for fluorescence imaging of tumors in patients. *Toxicol Rep.* 2019;6:409-15.
9. Boogerd LSF, Hoogstins CES, Schaap DP, Kusters M, Handgraaf HJM, van der Valk MJM, et al. Safety and effectiveness of SGM-101, a fluorescent antibody targeting carcinoembryonic antigen, for intraoperative detection of colorectal cancer: a dose-escalation pilot study. *The lancet Gastroenterology & hepatology.* 2018;3(3):181-91.
10. Hekman MC, Rijpkema M, Muselaers CH, Oosterwijk E, Hulsbergen-Van de Kaa CA, Boerman OC, et al. Tumor-targeted Dual-modality Imaging to Improve Intraoperative Visualization of Clear Cell Renal Cell Carcinoma: A First in Man Study. *Theranostics.* 2018;8(8):2161-70.
11. Hekman MCH, Rijpkema M, Bos DL, Oosterwijk E, Goldenberg DM, Mulders PFA, et al. Detection of Micrometastases Using SPECT/Fluorescence Dual-Modality Imaging in a CEA-Expressing Tumor Model. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine.* 2017;58(5):706-10.
12. Tummers WS, Warram JM, van den Berg NS, Miller SE, Swijnenburg RJ, Vahrmeijer AL, et al. Recommendations for reporting on emerging optical imaging agents to promote clinical approval. *Theranostics.* 2018;8(19):5336-47.
13. Lutje S, Rijpkema M, Helfrich W, Oyen WJ, Boerman OC. Targeted radionuclide and fluorescence dual-modality imaging of cancer: preclinical advances and clinical translation. *Molecular imaging and biology : MIB : the official publication of the Academy of Molecular Imaging.* 2014;16(6):747-55.
14. Lindmo T, Boven E, Cuttitta F, Fedorko J, Bunn PA, Jr. Determination of the immunoreactive fraction of radiolabeled monoclonal antibodies by linear extrapolation to binding at infinite antigen excess. *J Immunol Methods.* 1984;72(1):77-89.
15. Koppe MJ, Soede AC, Pels W, Oyen WJ, Goldenberg DM, Bleichrodt RP, et al. Experimental radioimmunotherapy of small peritoneal metastases of colorectal origin. *International journal of cancer.* 2003;106(6):965-72.
16. Rijpkema M, Oyen WJ, Bos D, Franssen GM, Goldenberg DM, Boerman OC. SPECT- and fluorescence image-guided surgery using a dual-labeled carcinoembryonic antigen-targeting antibody. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine.* 2014;55(9):1519-24.
17. Elekonawo FMK, de Gooyer JM, Bos DL, Goldenberg DM, Boerman OC, Brosens LAA, et al. Ex Vivo Assessment of Tumor-Targeting Fluorescent Tracers for Image-Guided Surgery. *Cancers (Basel).* 2020;12(4).
18. Kukis DL, DeNardo GL, DeNardo SJ, Mirick GR, Miers LA, Greiner DP, et al. Effect of the extent of chelate substitution on the immunoreactivity and biodistribution of 21T-BAT-Lym-1 immunoconjugates. *Cancer research.* 1995;55(4):878-84.
19. Al-Ejeh F, Darby JM, Thierry B, Brown MP. A simplified suite of methods to evaluate chelator conjugation of antibodies: effects on hydrodynamic radius and biodistribution. *Nucl Med Biol.* 2009;36(4):395-402.
20. Lewis MR, Raubitschek A, Shively JE. A facile, water-soluble method for modification of proteins with DOTA. Use of elevated temperature and optimized pH to achieve high specific activity and high chelate stability in radiolabeled immunoconjugates. *Bioconjugate chemistry.* 1994;5(6):565-76.
21. van Oudheusden TR, Braam HJ, Luyer MD, Wiezer MJ, van Ramshorst B, Nienhuijs SW, et al. Peritoneal cancer patients not suitable for cytoreductive surgery and HIPEC during explorative surgery: risk factors, treatment options, and prognosis. *Annals of surgical oncology.* 2015;22(4):1236-42.
22. Iversen LH, Rasmussen PC, Laurberg S. Value of laparoscopy before cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis. *The British journal of surgery.* 2013;100(2):285-92.

23. Pomel C, Appleyard TL, Gouy S, Rouzier R, Elias D. The role of laparoscopy to evaluate candidates for complete cytoreduction of peritoneal carcinomatosis and hyperthermic intraperitoneal chemotherapy. *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2005;31(5):540-3.
24. de Bree E, Koops W, Kroger R, van Ruth S, Witkamp AJ, Zoetmulder FA. Peritoneal carcinomatosis from colorectal or appendiceal origin: correlation of preoperative CT with intraoperative findings and evaluation of interobserver agreement. *Journal of surgical oncology*. 2004;86(2):64-73.
25. Schaap DP, de Valk KS, Deken MM, Meijer RPJ, Burggraaf J, Vahrmeijer AL, et al. Carcinoembryonic antigen-specific, fluorescent image-guided cytoreductive surgery with hyperthermic intraperitoneal chemotherapy for metastatic colorectal cancer. *The British journal of surgery*. 2020;107(4):334-7.
26. Brouwer NPM, Stijns RCH, Lemmens V, Nagtegaal ID, Beets-Tan RGH, Futterer JJ, et al. Clinical lymph node staging in colorectal cancer; a flip of the coin? *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2018;44(8):1241-6.
27. Boogerd LS, van der Valk MJ, Boonstra MC, Prevoo HA, Hilling DE, van de Velde CJ, et al. Biomarker expression in rectal cancer tissue before and after neoadjuvant therapy. *Onco Targets Ther*. 2018;11:1655-64.
28. de Gouw D, Rijpkema M, de Bitter TJJ, Baart VM, Sier CFM, Hernot S, et al. Identifying Biomarkers in Lymph Node Metastases of Esophageal Adenocarcinoma for Tumor-Targeted Imaging. *Mol Diagn Ther*. 2020.
29. van Hagen P, Hulshof MC, van Lanschot JJ, Steyerberg EW, van Berge Henegouwen MI, Wijnhoven BP, et al. Preoperative chemoradiotherapy for esophageal or junctional cancer. *The New England journal of medicine*. 2012;366(22):2074-84.
30. Vuijk FA, Hilling DE, Mieog JSD, Vahrmeijer AL. Fluorescent-guided surgery for sentinel lymph node detection in gastric cancer and carcinoembryonic antigen targeted fluorescent-guided surgery in colorectal and pancreatic cancer. *Journal of surgical oncology*. 2018;118(2):315-23.
31. Cuschieri A, Fayers P, Fielding J, Craven J, Bancewicz J, Joypaul V, et al. Postoperative morbidity and mortality after D1 and D2 resections for gastric cancer: preliminary results of the MRC randomised controlled surgical trial. The Surgical Cooperative Group. *Lancet (London, England)*. 1996;347(9007):995-9.
32. Bonenkamp JJ, Songun I, Hermans J, Sasako M, Welvaart K, Plukker JT, et al. Randomised comparison of morbidity after D1 and D2 dissection for gastric cancer in 996 Dutch patients. *Lancet (London, England)*. 1995;345(8952):745-8.
33. Hoogstins CES, Boogerd LSF, Sibinga Mulder BG, Mieog JSD, Swijnenburg RJ, van de Velde CJH, et al. Image-Guided Surgery in Patients with Pancreatic Cancer: First Results of a Clinical Trial Using SGM-101, a Novel Carcinoembryonic Antigen-Targeting, Near-Infrared Fluorescent Agent. *Annals of surgical oncology*. 2018;25(11):3350-7.
34. Hekman MC, Boerman OC, de Weijert M, Bos DL, Oosterwijk E, Langenhuijsen JF, et al. Targeted Dual-Modality Imaging in Renal Cell Carcinoma: An Ex Vivo Kidney Perfusion Study. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2016;22(18):4634-42.
35. Jay P, Berta P, Blache P. Expression of the carcinoembryonic antigen gene is inhibited by SOX9 in human colon carcinoma cells. *Cancer research*. 2005;65(6):2193-8.
36. Boonstra MC, de Geus SW, Prevoo HA, Hawinkels LJ, van de Velde CJ, Kuppen PJ, et al. Selecting Targets for Tumor Imaging: An Overview of Cancer-Associated Membrane Proteins. *Biomarkers in cancer*. 2016;8:119-33.

Figure 1: Biodistribution of 4 different dose levels of [^{111}In]-DTPA-SGM-101 in mice (n= 5 per dose level) with subcutaneous LS174T tumors. 1 group received a dose with an excess of 1 mg of unlabeled DTPA-SGM-101 as control group (gray, n= 5).

Figure 2: SPECT/CT imaging shows extensive metastatic disease within the abdominal cavity in a mouse with intraperitoneally induced tumors (A). After surgical exposure several tumor deposits are visible (B). Fluorescence imaging shows that after resection of macroscopically visible lesions several tumor deposits remain present within the abdominal cavity (C/D). After fluorescence-guided resection is performed, no additional lesions are visible within the abdominal cavity and only physiological uptake of the tracer in the liver remains (E).

Figure 3: Fluorescence-based (green, top graph) and radiosignal based (blue, bottom graph) tumor to background ratios after incubation of surgically obtained tissue specimens. Tumor deposits from case 9 and 10 were cut in half, each corresponding half was incubated with [^{111}In]-DTPA-SGM-101 (9* & 10*) or aspecific [^{111}In]-DTPA-IgG-Cy5.5 (9** & 10**). Case 7 was a CEA-negative malignancy (LAMN). Average tumor to background ratios are shown in orange.

Figure 4: Three examples of slides from the ex vivo incubation experiment on human tissues. From left to right: H&E staining, CEA staining, NIR-fluorescence and autoradiography. Clear co-localization of CEA-expressing tumor cells and both fluorescence and radiosignal is observed.

Figure 1




Figure 2




Figure 3





Figure 4



Clinical Cancer Research

Multimodal CEA-targeted image-guided colorectal cancer surgery using ¹¹¹In-labeled SGM-101

Jan Marie de Gooyer, Fortuné M.K. Elekonawo, Desirée L. Bos, et al.

Clin Cancer Res Published OnlineFirst September 8, 2020.

Updated version	Access the most recent version of this article at: doi: 10.1158/1078-0432.CCR-20-2255
Supplementary Material	Access the most recent supplemental material at: http://clincancerres.aacrjournals.org/content/suppl/2020/09/30/1078-0432.CCR-20-2255.DC1
Author Manuscript	Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://clincancerres.aacrjournals.org/content/early/2020/09/05/1078-0432.CCR-20-2255 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.